

Scalable Robust Principal Component Analysis using Grassmann Averages

— Supplementary Material —

Søren Hauberg, Aasa Feragen, Raffi Enficiaud, and Michael J. Black

APPENDIX A

We do not have explicit energy expressions for all the different variants of RGA, but in this appendix we consider the energy optimized by TGA(50%, 1) with unit weights. As this algorithm relies on trimmed spherical averages, we first consider these.

A.1 Trimmed Averages on S^{D-1}

In Euclidean spaces, the per-pixel trimmed average can be written as the solution to the following minimization problem

$$\boldsymbol{\mu}_{\text{Trim}, \mathbb{R}^D}(\mathbf{x}_{1:N}) = \arg \min_{\boldsymbol{\mu} \in \mathbb{R}^D} \sum_{n=1}^N \sum_{d=1}^D t_{nd} (x_{nd} - \mu_d)^2, \quad (33)$$

where the trimming weights $t_{nd} \in \{0, 1\}$ denote which elements are “trimmed away” and which are kept. For $P\%$ trimming we have

$$\bar{t} = \sum_{n=1}^N t_{nd} = N - \frac{2NP}{100} \quad \forall d = 1, \dots, D. \quad (34)$$

Note that \bar{t} is the same for all dimensions. The well-known solution to this problem is

$$\boldsymbol{\mu}_{\text{Trim}, \mathbb{R}^D}(\mathbf{x}_{1:N}) = \frac{1}{\bar{t}} \bar{\mathbf{x}}, \quad (35)$$

where $\bar{\mathbf{x}} \in \mathbb{R}^D$ has elements

$$\bar{x}_d = \sum_{n=1}^N t_{nd} x_{nd}. \quad (36)$$

- S. Hauberg is with the Section for Cognitive Systems at the Technical University of Denmark.
E-mail: sohau@dtu.dk
- A. Feragen is with the Department of Computer Science at the University of Copenhagen, Denmark.
E-mail: aasa@diku.dk
- R. Enficiaud is with the Software Workshop at the Max Planck Institute for Intelligent Systems in Tübingen, Germany.
E-mail: raffi.enficiaud@tue.mpg.de
- M.J. Black is with the Perceiving Systems Department at the Max Planck Institute for Intelligent Systems in Tübingen, Germany.
E-mail: black@tue.mpg.de

In the main paper, we consider extrinsic trimmed averages on the unit sphere. These can be similarly defined as

$$\begin{aligned} \boldsymbol{\mu}_{\text{Trim}, S^{D-1}}(\mathbf{u}_{1:N}) \\ = \arg \min_{\boldsymbol{\mu} \in S^{D-1}} \sum_{n=1}^N \sum_{d=1}^D t_{nd} (u_{nd} - \mu_d)^2. \end{aligned} \quad (37)$$

This constrained optimization problem has a simple closed-form solution:

Lemma 2. *Let $\mathbf{u}_n \in S^{D-1}$ and let the trimming weights t_{nd} be fixed, then Eq. 37 has solution*

$$\boldsymbol{\mu}_{\text{Trim}, S^{D-1}}(\mathbf{u}_{1:N}) = \frac{1}{\|\bar{\mathbf{u}}\|} \bar{\mathbf{u}}, \quad (38)$$

where $\bar{\mathbf{u}} = \sum_{n=1}^N t_{nd} \mathbf{u}_{nd}$ is the Euclidean trimmed average of the \mathbf{u}_{nd} .

Proof: The results follows by straight-forward computations: We seek the minima of

$$f(\boldsymbol{\mu}) = \sum_{n=1}^N \sum_{d=1}^D t_{nd} (u_{nd} - \mu_d)^2 \quad (39)$$

subject to the constraint $\|\boldsymbol{\mu}\| = 1$. We write the constraints using a Lagrange-multiplier

$$\hat{f}(\boldsymbol{\mu}, \lambda) = f(\boldsymbol{\mu}) + \lambda g(\boldsymbol{\mu}), \quad \text{where} \quad (40)$$

$$g(\boldsymbol{\mu}) = 1 - \boldsymbol{\mu}^\top \boldsymbol{\mu} = 1 - \sum_{d=1}^D \mu_d^2. \quad (41)$$

We evaluate derivatives as

$$\frac{\partial f}{\partial \mu_d} = -2 \sum_{n=1}^N t_{nd} (u_{nd} - \mu_d) \quad (42)$$

$$= -2 \sum_{n=1}^N t_{nd} u_{nd} + 2 \sum_{n=1}^N t_{nd} \mu_d \quad (43)$$

$$= -2 (\bar{u}_d - \bar{t} \mu_d). \quad (44)$$

$$\frac{\partial g}{\partial \mu_d} = -2 \mu_d \quad (45)$$

$$\frac{\partial \hat{f}}{\partial \mu_d} = -2 (\bar{u}_d - \bar{t} \mu_d) - 2 \lambda \mu_d \quad (46)$$

$$= -2 (\bar{u}_d - (\bar{t} - \lambda) \mu_d). \quad (47)$$

Setting $\partial \hat{f} / \partial \mu_d = 0$ gives

$$\mu_d = \frac{\bar{u}_d}{\bar{t} - \lambda}. \quad (48)$$

We evaluate λ by setting $\partial \hat{f} / \partial \lambda = 0$:

$$\frac{\partial \hat{f}}{\partial \lambda} = 1 - \sum_{d=1}^D \mu_d^2 = 1 - \sum_{d=1}^D \frac{\bar{u}_d^2}{(\bar{t} - \lambda)^2} \quad (49)$$

$$= 1 - (\bar{t} - \lambda)^{-2} \|\bar{\mathbf{u}}\|^2 = 0 \Rightarrow \quad (50)$$

$$\bar{t} - \lambda = \pm \|\bar{\mathbf{u}}\|. \quad (51)$$

Combining Eq. 48 and 51 gives

$$\mu_d = \pm \frac{1}{\|\bar{\mathbf{u}}\|} \bar{u}_d. \quad (52)$$

The unknown sign is determined by evaluating f at both choices and picking the smaller option. \square

The per-pixel trimmed spherical average, thus, has the closed-form solution given by the per-pixel trimmed Euclidean average projected onto the sphere. In the case of 50% trimming, the per-pixel trimmed average coincides with the per-pixel median, and it follows that

$$\boldsymbol{\mu}_{\text{Median}, S^{D-1}}(\mathbf{u}_{1:N}) = \arg \min_{\boldsymbol{\mu} \in S^{D-1}} \sum_{n=1}^N \sum_{d=1}^D |u_{nd} - \mu_d| \quad (53)$$

can be solved by the per-pixel median projected onto the unit-sphere.

A.2 The Energy Optimized by TGA(50%, 1)

Intuitively, TGA(50%, 1) should find the ‘‘median subspace’’ spanned by the data. Indeed it optimizes the energy

$$\boldsymbol{\mu}_{\text{Median}}(\mathbf{u}_{1:N}) = \arg \min_{\boldsymbol{\mu} \in [\boldsymbol{\mu}]} \left\{ \sum_{n=1}^N \sum_{d=1}^D |u_{nd} - \mu_d| \right\}, \quad (54)$$

which can be interpreted as a pixel-wise median subspace.

At every step, the TGA algorithm updates the representatives \mathbf{u}_n of the equivalence class $[\mathbf{u}_n]$, to $\alpha_n \mathbf{u}_n$ for some element of the antipodal group $\alpha_n \in \{\pm 1\}$. To obtain a convergence guarantee, we assume that the selection of the signs α_n are made to optimize the chordal L_1 distance to the current mean estimate, that is¹

$$\alpha_n = \arg \min_{a_n = \pm 1} \sum_{d=1}^D |a_n u_{nd} - \mu_d|. \quad (55)$$

Lemma 3. *Then, with probability 1, TGA(50%, 1) converges to a local minimum of Eq. 54 in finite time.*

Proof: We shall show that, with probability 1, there exists $M \in \mathbb{N}_0$ such that $\alpha_n = 1$ for all n in every iteration after the M^{th} iteration of the algorithm. Moreover, the value of the energy function

$$\sum_{n=1}^N \sum_{d=1}^D |u_{nd} - \mu_d| \quad (56)$$

1. In practical implementations we pick α_n to optimize the L_2 distance rather than the L_1 as this can be done highly efficiently.

decreases strictly for steps 1 to $(M - 1)$, that is

$$\sum_{n=1}^N \sum_{d=1}^D |\alpha_n u_{nd} - \mu_d| < \sum_{n=1}^N \sum_{d=1}^D |u_{nd} - \mu_d| \quad (57)$$

for every iteration up to M .

In the i^{th} iteration, with probability 1, we have

$$\sum_{d=1}^D |-1 \cdot u_{nd} - \mu_d| \neq \sum_{d=1}^D |u_{nd} - \mu_d| \quad (58)$$

for every $n = 1 \dots N$, because the set on which $\sum_{d=1}^D |-1 \cdot u_{nd} - \mu_d| = \sum_{n=1}^N \sum_{d=1}^D |u_{nd} - \mu_d|$ has measure 0.

Now, we could have $\alpha_n = 1$ for all n , in which case the algorithm has converged and $i \geq M$. Otherwise, there exists some n for which $\alpha_n = -1$ gives

$$\sum_{d=1}^D |\alpha_n \cdot u_{nd} - \mu_d| < \sum_{d=1}^D |u_{nd} - \mu_d|, \quad (59)$$

in which case the energy in Eq. 56 will decrease strictly in the i^{th} iteration.

The fact that M exists and the TGA algorithm converges in a finite number of steps follows from the fact that there are only finitely many ways to change the sign of $\mathbf{u}_{1:N}$, each giving a fixed value of the energy function (56), so there cannot be an infinite sequence of strictly decreasing values.

As a very small perturbation of the data points will not lead to a change in the signs α_n , the algorithm must moreover converge to a local optimum. \square